arXiv:math/0603667v1 [math.GR] 28 Mar 2006

On reflections in Coxeter groups commuting with a reflection

Koji Nuida

Abstract

We consider the subgroup of an arbitrary Coxeter group generated by the reflections which commute with a given reflection. We determine completely for which case this subgroup is finitely generated. This result provides new uniform examples of non-finitely generated subgroups in finitely generated groups.

1 Introduction

Let (W, S) be an arbitrary Coxeter system, i.e. W is a Coxeter group and S is the set of Coxeter generators. We call an element $w \in W$ a reflection in W if and only if w is conjugate to some element of S. When realizing W canonically as a reflection group on a real vector space V (the geometric representation space) equipped with (not necessarily nondegenerate) symmetric bilinear form, this definition is equivalent to that w is a reflection in the space V geometrically. A subgroup of W generated by some reflections is called a reflection subgroup. Thus reflection subgroups are one of the most important subgroups of Coxeter groups, both group-theoretically and geometrically. For example, a theorem of Vinay V. Deodhar [4] or Matthew Dyer [5, Theorem 3.3] proves that any reflection subgroup of a Coxeter group is also a Coxeter group.

In this paper, we treat the following reflection subgroup $W^{\perp t}$ of an arbitrary Coxeter group W. This is generated by the reflections, other than a given reflection $t \in W$, which commutes with t (or equivalently, whose -1-eigenvector is perpendicular to that of t). The aim of this paper is to determine which subgroups $W^{\perp t}$ are finitely generated (Theorems 4.13 and 4.14). Indeed, we consider the case $t \in S$ only, since a general case can be easily reduced to this special case. On the other hand, although the most interesting case would be the one that W is finitely generated (for which our result is summarized as Corollary 4.15), we decide to treat an arbitrary W, since the argument does not become too long and complicated.

One of the most significant phenomena in group theory is the existence of non-finitely generated subgroups in finitely generated groups. Corollary 4.15 says that even such a naturally and uniformly introduced reflection subgroup $W^{\perp t}$ of a finitely generated Coxeter group W is frequently non-finitely generated. We hope that our result enriches the list of interesting examples of the significant phenomena. (Note that some other uniform examples are also provided in a recent paper [7] of Swiatoslaw R. Gal; see [7, Proposition 2.1 and Remark 2.2].)

This paper is organized as follows. Section 2 is a preliminary for graphs, groupoids and Coxeter groups. Our argument here is based on the preceding results by Brigitte Brink [2] and by the author [11], which describe the structure of $W^{\perp t}$ as a Coxeter group combinatorially (note that there have been many other results describing the structure; see e.g. [8, 1, 3]). Section 3 summarizes these preceding results and gives some further remarks. Finally, Section 4 gives our main results (Theorems 4.13 and 4.14), together with some corollaries (Corollaries 4.15–4.19) which are obtained by restricting the main theorems to some subclasses of Coxeter groups.

Acknowledgement. The author would like to express his deep gratitude to Professor Itaru Terada and to Professor Kazuhiko Koike for their precious advice and encouragement. Moreover, the author was supported by JSPS Research Fellowship throughout this research.

2 Preliminaries

In this paper, we abbreviate the notation $\{x\}$ for a set with a single element to x, unless some ambiguity occurs.

2.1 Graphs

All graphs appearing in this paper are unoriented and simple, i.e. having no loops and no multiple edges. So any path in a graph G is denoted, as usual, by a sequence of its vertices. In particular, the path of length 0 with starting point x is denoted by (x). Let $V(G)$ denote the vertex set of a graph G, which is not assumed to be finite. A *cycle* in G signifies a closed path $(x_0, x_1, \dots, x_{n-1}, x_n)$ with no repetitions of vertices except $x_0 = x_n$ and no shortcutting edges (x_i, x_j) in G such that $i - j \not\equiv 0, \pm 1 \pmod{n}$. For $x \in V(G)$, let $G_{\sim x}$ be the connected component of G containing x. For $I \subseteq V(G)$, let $G|_I$ denote the restriction of G with vertex set I.

Definition 2.1. *For a graph G, we refer to a subset $I \subseteq V(G)$ as a* pre-cycle core *of G if $G|_I$ is connected and contains all cycles in G. The unique minimal pre-cycle core of G (if exists) is referred to as the* cycle core *of G.*

Here we introduce a decomposition of a connected graph associated to its pre-cycle core.

Lemma 2.2. *Let G be a connected graph and $I \subseteq V(G)$ its pre-cycle core. Then G is the union of the subgraph $G|_I$ and mutually disjoint subtrees T_x, indexed by $x \in I$, such that $V(T_x) \cap I = \{x\}$ for any $x \in I$.*

Proof. First we show that for each $s \in V(G)$, there exists a unique vertex $x_s \in I$ such that some path in G from s to x_s contains no other vertex in I. Since G is connected, the existence is obvious, so we consider the uniqueness assertion. This is obvious if $s \in I$ (now $x_s = s$), so suppose not. Now assume contrary that two such vertices $y \neq z$ in I exist. Then these are joined by a path in G which does not intersect $G|_I$ except at the endpoints y and z. Since $G|_I$ is connected by the hypothesis, this implies that G contains a cycle which is not contained in $G|_I$, contradicting the choice of I. Thus the uniqueness also holds.

Now our claim holds by taking $T_x = G|_{J_x}$ for each $x \in I$, where $J_x = \{s \in V(G) \mid x_s = x\}$. Note that T_x is connected and $J_x \cap I = \{x\}$ by definition, so T_x is a subtree of G since I is a pre-cycle core. The existence and the uniqueness of x_s for each $s \in V(G)$ show, respectively, that $V(G) = I \cup \bigcup_{x \in I} V(T_x)$ and all T_x are disjoint. Moreover, the uniqueness also implies that two vertices $s \in J_x$ and $t \in J_y$, with $x, y \in I$ distinct, cannot be joined by an edge of G, unless $s = x$ and $t = y$. Hence G is the union of $G|_I$ and all T_x, as desired. $\square$

By the shape of the decomposition, we have the following immediately.

Corollary 2.3. *Let G be a connected graph and $I \subseteq V(G)$ its pre-cycle core. Then for any $y, z \in I$, every non-backtracking path in G between y and z is contained in $G|_I$.*

Regarding the cycle core, the next lemma follows from Corollary 2.3.

Lemma 2.4. *Suppose that a graph G is connected and contains a cycle. Then the intersection of (possibly infinitely many) pre-cycle cores of G is also a pre-cycle core of G. Hence the cycle core of such G exists.*

Proof. Let $\{I_\lambda\}_\lambda$ be a family of pre-cycle cores of G and $I = \bigcap_\lambda I_\lambda$. Then any cycle in G is contained in all $G|_{I_\lambda}$, so in $G|_I = \bigcap_\lambda G|_{I_\lambda}$ as well. Moreover, if $y, z \in I$, then a non-backtracking path in G between y and z (which exists since G is connected) is contained in all $G|_{I_\lambda}$ by Corollary 2.3 (note that $y, z \in I \subseteq I_\lambda$), so in $G|_I$ as well. This means that $G|_I$ is also connected. Hence the claim holds. □

2.2 Groupoids

A *groupoid* is a small category whose morphisms are all invertible; namely a family of sets $\mathcal{G} = \{\mathcal{G}_{x,y}\}_{x,y \in V(\mathcal{G})}$ with some index set (or *vertex set*) $V(\mathcal{G})$ endowed with (1) multiplications $\mathcal{G}_{x,y} \times \mathcal{G}_{y,z} \to \mathcal{G}_{x,z}$ satisfying the associativity law, (2) an identity element $1 = 1_x$ in each $\mathcal{G}_{x,x}$, and (3) an inverse $g^{-1} \in \mathcal{G}_{y,x}$ for every $g \in \mathcal{G}_{x,y}$. Each $\mathcal{G}_{x,x}$ is a group, called a *vertex group* of $\mathcal{G}$ and denoted by $\mathcal{G}_x$. A *homomorphism* between groupoids is a covariant functor between them regarded as categories. Notions such as *isomorphisms* and *subgroupoids* are defined as usual.

The *fundamental groupoid* of a graph G, denoted by $\pi_1(G; *, *)$ in this paper, is a groupoid with vertex set $V(G)$. The edges of G, endowed with orientation from a vertex x to another y, define elements of $\pi_1(G; x, y) = \pi_1(G; *, *)_{x,y}$; and $\pi_1(G; *, *)$ is freely generated by these elements. So a path $(x_0, x_1, \dots, x_n)$ in G represents an element of $\pi_1(G; x_0, x_n)$. The same edge or path endowed with two opposite orientations give inverse elements to each other. Each element of $\pi_1(G; *, *)$ is represented by a unique non-backtracking path. A vertex group $\pi_1(G; x) = \pi_1(G; x, x)$ is the *fundamental group* of G at x, which is a free group. This coincides with the usual fundamental group of G when regarded as a 1-cell complex. Note that, if H is a subgraph of G, then $\pi_1(H; *, *)$ is naturally a subgroupoid of $\pi_1(G; *, *)$.

Lemma 2.5. *Let G be a connected graph and $I \subseteq V(G)$.*

1. *If I is a pre-cycle core of G, then $\pi_1(G|_I; x, y) = \pi_1(G; x, y)$ for any $x, y \in I$.*
2. *Conversely, if $x \in I$ and $\pi_1(G|_I; x) = \pi_1(G; x)$, then the connected component $(G|_I)_{\sim x}$ contains all cycles in G.*

Proof. Claim 1 follows from Corollary 2.3, since each element of $\pi_1(G; x, y)$ is represented by a non-backtracking path between x and y. For Claim 2, we may assume that $G|_I$ is connected. Then for any cycle C in G, since G is connected, there is a non-backtracking closed path P in G starting from x and containing C. Now $P \in \pi_1(G|_I; x)$ by the hypothesis, so it follows that $C \subseteq G|_I$. Hence the claim holds. □

2.3 Coxeter groups

In this paper, let W denote a Coxeter group, S a specified (possibly infinite) generating set of W (so (W, S) is a Coxeter system), and $m_{s,t} = m(s, t)$ the (possibly infinite) order of the product st of two generators $s, t \in S$ in W. Thus $m_{s,t} = m_{t,s}$ and $m_{s,s} = 1$ for any $s, t \in S$, and W admits the following group presentation

$$W = \langle S \mid (st)^{m_{s,t}} = 1 \text{ if } s, t \in S \text{ and } m_{s,t} < \infty \rangle.$$

We refer to the book [9] for basics of Coxeter groups, unless otherwise noticed. An *isomorphism of Coxeter systems* $(W, S) \to (W', S')$ signifies a group isomorphism $W \to W'$ mapping S onto S'. Let Γ denote the Coxeter graph of W, which is a graph with vertex set S and in which two vertices $s, t \in S$ are joined by an edge with label $m_{s,t}$ if and only if $3 \leq m_{s,t} \leq \infty$ (this label is usually omitted when $m_{s,t} = 3$). For $w \in W$, let $\ell(w)$ be the smallest integer r such that $w = s_1 s_2 \cdots s_r$ for some $s_i \in S$ (the *length* of w). The map $\ell : W \to \{0, 1, 2, \dots\}$ is called the *length function* of W.

Definition 2.6. *For an expression $w = s_1 s_2 \cdots s_r$ of $w \in W$ with $r = \ell(w)$, define* $\mathrm{supp}(w) = \{s_1, s_2, \dots, s_r\} \subseteq S$, *the* support *of w.*

It is well known that the support of $w \in W$ is independent of the choice of the expression. This yields immediately the following property:

Lemma 2.7. *Let $w_i \in W$ $(1 \leq i \leq k)$, $w = w_1 w_2 \cdots w_k$ and suppose that $\ell(w) = \sum_{i=1}^{k} \ell(w_i)$. Then* $\mathrm{supp}(w) = \bigcup_{i=1}^{k} \mathrm{supp}(w_i)$.

On the other hand, since no proper subset of S generates W, the following property holds:

Proposition 2.8. *A Coxeter group W is finitely generated if and only if the generating set S is finite.*

Proof. The 'if' part is trivial. For the "only if" part, suppose that W has a finite generating set X. Then the set $I = \bigcup_{w \in X} \mathrm{supp}(w)$ is also finite and generates W, so $I = S$, therefore $|S| < \infty$ as desired. □

For $I \subseteq S$, let W_I denote the subgroup of W generated by I, called a (*standard*) *parabolic subgroup*. It is fundamental that (W_I, I) is a Coxeter system with Coxeter graph $\Gamma_I = \Gamma|_I$ and length function $\ell_I = \ell|_{W_I}$, and $\bigcap_\lambda W_{J_\lambda} = W_J$ (where $J = \bigcap_\lambda J_\lambda$) for a family $\{J_\lambda\}_\lambda$ of subsets of S. We say that the graph Γ_I is of type $\widetilde{A}_{n-1}$ if Γ_I is a cycle with n vertices and all edges of Γ_I are labelled 3 (or, in our convention, are not labelled). A parabolic subgroup W_I, with I the vertex set of a connected component of Γ, is called an *irreducible component* of W; and W is called *irreducible* if Γ is connected. Now W is the (restricted) direct product of the irreducible components.

Let V denote the *geometric representation space* (over $\mathbb{R}$) endowed with basis $\Pi = \{\alpha_s \mid s \in S\}$ and symmetric bilinear form $\langle\,,\,\rangle$ determined by

$$\langle \alpha_s, \alpha_t \rangle = \begin{cases} -\cos(\pi/m_{s,t}) & \text{if } m_{s,t} < \infty; \\ -1 & \text{if } m_{s,t} = \infty. \end{cases}$$

Then $\langle\,,\,\rangle$ is W-invariant, where W acts on V by $s \cdot v = v - 2\langle \alpha_s, v \rangle \alpha_s$ for $s \in S$ and $v \in V$. Let $\Phi = W \cdot \Pi$, $\Phi^+ = \Phi \cap \mathbb{R}_{\geq 0}\Pi$ and $\Phi^- = -\Phi^+$ denote the *root system*, the set of *positive roots* and the set of *negative roots*, respectively. Each element of Φ (a *root*) is a unit vector with respect to $\langle\,,\,\rangle$, and $\Phi = \Phi^+ \sqcup \Phi^-$. Write $\Psi^+ = \Psi \cap \Phi^+$ for any subset $\Psi \subseteq \Phi$. For a vector $v = \sum_{s \in S} c_s \alpha_s \in V$, define

$$\mathrm{supp}(v) = \{s \in S \mid c_s \neq 0\}, \text{ the } \textit{support} \text{ of } v.$$

For a subset $I \subseteq S$, let V_I denote the subspace of V spanned by the set $\Pi_I = \{\alpha_s \mid s \in I\}$, and put $\Phi_I = \Phi \cap V_I$. It is well known that

$$\Phi_I = W_I \cdot \Pi_I,$$

the root system of (W_I, I) (see e.g. [6, Lemma 4]).

For a root $\gamma = w \cdot \alpha_s \in \Phi$, let s_γ denote the *reflection* $wsw^{-1} \in W$ along the root γ, acting on V by $s_\gamma \cdot v = v - 2\langle \gamma, v \rangle \gamma$ for $v \in V$. Then $w \in W$ is a reflection (along some root) if and only if w is conjugate to an element of S. Moreover, it is known that $w \in W$ is a reflection in this sense if and only if w is geometrically a reflection in the vector space V. A subgroup of W generated by some reflections is called a *reflection subgroup*.

The following lemma will be used later.

Lemma 2.9 ([10, Lemma 2.7]). *Let $1 \neq w \in W$, $I = \mathrm{supp}(w)$, $\gamma \in \Phi^+$ and $J = \mathrm{supp}(\gamma)$. Suppose that $I \cap J = \emptyset$ and J is adjacent to I in the Coxeter graph Γ. Then $w \cdot \gamma \in \Phi_{I \cup J}^+ \smallsetminus \Phi_J$, hence $w \cdot \gamma \neq \gamma$.*

The following subgraph of a Coxeter graph plays an important role later.

Definition 2.10. *For a subset $I \subseteq S$, we define the* odd Coxeter graph Γ_I^{odd} *to be the subgraph of Γ_I obtained by removing all edges with labels either even or ∞. Moreover, for $s \in I$, let $I_{\sim s}^{\mathrm{odd}}$ denote the vertex set of the connected component $(\Gamma_I^{\mathrm{odd}})_{\sim s}$ of Γ_I^{odd} containing s.*

At last of this subsection, we prepare one more notation.

Definition 2.11. *For any subsets $I \subseteq J \subseteq S$, let $J_{\sim I}$ denote the union of vertex sets of connected components of Γ_J containing some vertex in I.*

3 Note on centralizers of reflections

In a paper [2], Brigitte Brink showed that the centralizer $Z_{W_I}(x)$ of a generator $x \in I$ in an arbitrary parabolic subgroup W_I admits the following decomposition

$$Z_{W_I}(x) = (\langle x \rangle \times W_I^{\perp x}) \rtimes Y_x^I = \langle x \rangle \times (W_I^{\perp x} \rtimes Y_x^I)$$

(where the notations follow the author's paper [11], which enhances the result in [2] in more general setting, with slight modification). The main object of the present paper is the subgroup $W_I^{\perp x}$, which is generated by the reflections along the roots in the following set

$$\Phi_I^{\perp x} = \{\gamma \in \Phi_I \mid \langle \gamma, \alpha_x \rangle = 0\}.$$

Let Π_I^x be the "simple system" in $(\Phi_I^{\perp x})^+$, i.e. the members of Π_I^x are the roots in $(\Phi_I^{\perp x})^+$ which is not a positive linear combination of at least two roots in $(\Phi_I^{\perp x})^+$. Let

$$R_I^x = \{s_\gamma \mid \gamma \in \Pi_I^x\}$$

be the set of corresponding reflections. Then $(W_I^{\perp x}, R_I^x)$ is a Coxeter system by a theorem of Vinay V. Deodhar [4] or Matthew Dyer [5, Theorem 3.3]. In [11], the precise structure of $W_I^{\perp x}$ is described by using the group Y_x^I.

For the group Y_x^I, Brink proved in [2] the following result, where we put

$$\mathcal{F}^I = \pi_1(\Gamma_I^{\mathrm{odd}}; *, *) \text{ (see Section 2.2 and Definition 2.10 for notations)}$$

and Y^I is the groupoid with vertex set I defined by

$$Y_{y,z}^I = \{w \in W_I \mid \alpha_y = w \cdot \alpha_z \text{ and } w \cdot (\Phi_I^{\perp z})^+ \subseteq \Phi^+\} \text{ for } y, z \in I$$

(note that Y^I is indeed a groupoid; see [2] or [11]).

Theorem 3.1. *There is a unique groupoid isomorphism $\pi^I : \mathcal{F}^I \to Y^I$ such that $\pi^I(y) = y$ for $y \in I$,*

$$\pi^I(y, z) = (zy)^k \in Y_{y,z}^I \text{ for } y, z \in I \text{ with } m_{y,z} = 2k+1 \text{ odd}$$

and $\ell(\pi^I(y_0, y_1, \dots, y_n)) = \sum_{i=1}^n \ell(\pi^I(y_{i-1}, y_i))$ if $(y_0, y_1, \dots, y_n)$ is a non-backtracking path in Γ_I^{odd}.

Thus Y_x^I is isomorphic via π^I to the free group $\mathcal{F}_x^I = \pi_1(\Gamma_I^{\mathrm{odd}}; x)$.

Now we come back to the description of $W_I^{\perp x}$. First note that, if $y, z \in I$ are distinct and (when $m_{y,z} < \infty$) w is the longest element of $W_{\{y,z\}}$ (i.e. $w = (zy)^k$ if $m_{y,z} = 2k$ is even, and $w = (yz)^k y$ if $m_{y,z} = 2k+1$ is odd), then we have $|W_{\{y,z\}_{\sim z}}| < \infty$ (see Definition 2.11 for notation) and $w_{y,z} y w_{y,z}^{-1} = y$ (where $w_{y,z} = wy$) if and only if $m_{y,z}$ is even. By this observation, the results in [11] (especially Example 4.12 and Theorem 4.13) imply, when restricted to this case, that the set R_I^x consists of the elements $w \pi^I(p^I_{x,y}) w_{y,z} \pi^I(p^I_{y,x}) w^{-1}$ (which is denoted in [11] by $r_x(w, w_{y,z})$) with $w \in Y_x^I$ and $(y,z) \in \mathcal{E}_x^I$, where

$$\mathcal{E}_x^I = \{(s,t) \mid s \in I^{\mathrm{odd}}_{\sim x},\ t \in I \smallsetminus s \text{ and } m_{s,t} \text{ is even}\}$$

(see Definition 2.10 for notation) and $p^I_{s,t}$ (with $s,t \in I^{\mathrm{odd}}_{\sim x}$) is the unique non-backtracking path between s and t in a fixed maximal tree $\mathcal{T}$ in the connected component $(\Gamma_I^{\mathrm{odd}})_{\sim x}$ of Γ_I^{odd} containing x; so that in the groupoid $\mathcal{F}^I$, we have $p^I_{s,t} \in \mathcal{F}^I_{s,t}$, $p^I_{s,s} = (s)$, $p^I_{t,s} = (p^I_{s,t})^{-1}$ and $p^I_{s,s'} \cdot p^I_{s',t} = p^I_{s,t}$.

Now we introduce a new notation $r_x^I(y,z;c)$ for an element $r_x(w, w_{y,z})$ in the original notation, where $c \in \mathcal{F}^I_{x,y}$ and $\pi^I(c) = w \pi^I(p^I_{x,y})$; namely

$$r_x^I(y,z;c) = \pi^I(c) w_{y,z} \pi^I(c)^{-1} \text{ for } (y,z) \in \mathcal{E}_x^I \text{ and } c \in \mathcal{F}^I_{x,y}.$$

Write $\xi = (\xi^o, \xi^\dagger)$ for each $\xi \in \mathcal{E}_x^I$. Moreover, if $\xi \in \mathcal{E}_x^I$ and we put

$$\widetilde{\alpha}_{\xi^o,\xi^\dagger} = \frac{1}{\sin(\pi/m)} (\cos(\pi/m) \alpha_{\xi^o} + \alpha_{\xi^\dagger}) \text{ where } m = m_{\xi^o,\xi^\dagger} \text{ is even,}$$

then the element $w_{\xi^o,\xi^\dagger} = (\xi^\dagger \xi^o)^{k-1} \xi^\dagger$, where $m = 2k$, is the reflection along the root $\widetilde{\alpha}_{\xi^o,\xi^\dagger} \in (\Phi^{\perp \xi^o}_{\{\xi^o,\xi^\dagger\}})^+$ (note that $w_{\xi^o,\xi^\dagger}$ is a reflection by the shape of the expression, and a vector $v \in V_{\{\xi^o,\xi^\dagger\}}$ is uniquely determined by the conditions that the coefficients in v are nonnegative, $\langle v, v \rangle = 1$ and $\langle v, \alpha_{\xi^o} \rangle = 0$). Thus $r_x^I(\xi;c)$ is the reflection along the root

$$\gamma_x^I(\xi;c) = \pi^I(c) \cdot \widetilde{\alpha}_{\xi^o,\xi^\dagger},$$

so we have $\Pi_I^x = \{\gamma_x^I(\xi;c) \mid \xi \in \mathcal{E}_x^I \text{ and } c \in \mathcal{F}^I_{x,\xi^o}\}$. We write $r_x^I(\xi) = r_x^I(\xi; p^I_{x,\xi^o})$ and $\gamma_x^I(\xi) = \gamma_x^I(\xi; p^I_{x,\xi^o})$ for simplicity.

We consider relations between the elements of R_I^x. For $\xi \in \mathcal{E}_x^I$ and $z \in I \smallsetminus \{\xi^o, \xi^\dagger\}$ such that $J_{\sim\{\xi^o,\xi^\dagger\}}$ generates a finite parabolic subgroup where $J = \{\xi^o, \xi^\dagger, z\}$ (see Figure 1 for classification of such cases), there is a unique $\zeta \in \mathcal{E}_x^I \smallsetminus \xi$ such that $\zeta^o, \zeta^\dagger \in J$ and the graph $(\Gamma_J^{\mathrm{odd}})_{\sim \xi^o}$ is a path (say q) between ξ^o and ζ^o, where q is oriented as satisfying that $q \in \mathcal{F}^I_{\xi^o,\zeta^o}$. Then we introduce a relation $(\xi;c) \overset{k}{\sim}_I (\zeta; cq)$ for each $c \in \mathcal{F}^I_{x,\xi^o}$, where k is the (finite) order of the product of $r_x^I(\xi) = r_x^I(\xi; p^I_{x,\xi^o})$ and $r_x^I(\zeta; p^I_{x,\xi^o} q)$. See Figure 1 for precise values of the k; see also [11, Tables 1–2]. These relations are in fact symmetric; let $\sim_I$ denote the transitive closure of $\overset{1}{\sim}_I$. Now the next theorem is a special case of [11, Theorem 4.14].

Theorem 3.2. *The pair* $(W_I^{\perp x}, R_I^x)$ *is a Coxeter system, and* R_I^x *consists of the elements* $r_x^I(\xi; c)$ *with* $\xi \in \mathcal{E}_x^I$ *and* $c \in \mathcal{F}_{x,\xi^o}^I$ *defined above. Moreover, for* $\xi, \zeta \in \mathcal{E}_x^I$*,* $c \in \mathcal{F}_{x,\xi^o}^I$*,* $d \in \mathcal{F}_{x,\zeta^o}^I$ *and* $2 \leq m < \infty$*,*

1. *we have* $r_x^I(\xi; c) = r_x^I(\zeta; d)$ *if and only if* $(\xi; c) \sim_I (\zeta; d)$*;*

2. *the product of* $r_x^I(\xi; c)$ *and* $r_x^I(\zeta; d)$ *has order* m *if and only if* $(\xi; c) \sim_I (\xi'; c') \overset{m}{\sim}_I (\zeta'; d') \sim_I (\zeta; d)$ *for some pairs* $(\xi'; c')$ *and* $(\zeta'; d')$*.*

Here we give some graphical explanation for the relation $\overset{1}{\sim}_I$. We regard a relation $(\xi; c) \overset{1}{\sim}_I (\zeta; cq)$ as *moves* in I of two particles from the vertices ξ^o and $\xi^\dagger$ to the vertices ζ^o and $\zeta^\dagger$, respectively, where the particle at ξ^o (which we refer to as the *mover* of ξ) moves to ζ^o along the path q in Γ_I^{odd}. (We refer to the other particle as the *support* of ξ.) So the path q is the trace of the mover under this move. Now the moves are classified into two types depicted in Figures 2 and 3; we call the first one a *sliding*; and the second one a *switching* (at a vertex z). Note that all the moves are invertible. On the other hand, the trace $\widetilde{q}$ of the mover under some consecutive moves in I from ξ to ζ is a path in Γ_I^{odd} between ξ^o and ζ^o; now we say that the element $\widetilde{q} \in \mathcal{F}_{\xi^o,\zeta^o}^I$ is *realized* by these consecutive moves. Using the terminology, Theorem 3.2 (1) is restated as follows:

Theorem 3.3. *For* $\xi, \zeta \in \mathcal{E}_x^I$*,* $c \in \mathcal{F}_{x,\xi^o}^I$ *and* $d \in \mathcal{F}_{x,\zeta^o}^I$*, we have* $r_x^I(\xi; c) = r_x^I(\zeta; d)$ *if and only if there are consecutive moves in* I *from* ξ *to* ζ *which realize the element* $c^{-1}d \in \mathcal{F}_{\xi^o,\zeta^o}^I$*.*

Example 3.4. *In this and the next examples, let* $x \in I \subseteq S$ *such that* Γ_I^{odd} *is a cycle* $P = (y_0, y_1, \dots, y_{n-1}, y_n = y_0)$ *with* $n \geq 4$*, and put* $\xi_{i,j} = (y_i, y_j)$ *and* $m_{i,j} = m_{y_i,y_j}$*.*

In this example, suppose that $n = 2N + 1$ *is odd (so* $N \geq 2$*) and* $m_{0,k}$ *is even for some* $2 \leq k \leq n - 2$*. Then by definition of the moves, if some nonempty non-backtracking sequence of consecutive moves in* I *from* $\xi_{k,0}$ *to* $\xi_{k,0}$ *exists, then the shortest one is (up to orientation) the following:*

$$\begin{aligned}
&\xi_{k,0} \to \xi_{k+1,0} \to \xi_{k+2,0} \to \cdots \to \xi_{2N-1,0} \overset{2N}{\rightsquigarrow} \xi_{0,2N-1} \\
&\to \xi_{1,2N-1} \to \cdots \to \xi_{2N-3,2N-1} \overset{2N-2}{\rightsquigarrow} \xi_{2N-1,2N-3} \to \xi_{2N,2N-3} \\
&\to \cdots \to \xi_{2N-5,2N-3} \overset{2N-4}{\rightsquigarrow} \xi_{2N-3,2N-5} \to \cdots \to \xi_{2N-7,2N-5} \overset{2N-6}{\rightsquigarrow} \xi_{2N-5,2N-7} \\
&\to \cdots\cdots \to \xi_{1,3} \overset{2}{\rightsquigarrow} \xi_{3,1} \to \cdots \to \xi_{2N,1} \overset{0}{\rightsquigarrow} \xi_{1,2N} \\
&\to \cdots \to \xi_{2N-2,2N} \overset{2N-1}{\rightsquigarrow} \xi_{2N,2N-2} \to \cdots \to \xi_{2N-4,2N-2} \overset{2N-3}{\rightsquigarrow} \xi_{2N-2,2N-4} \\
&\to \cdots\cdots \to \xi_{2,4} \overset{3}{\rightsquigarrow} \xi_{4,2} \to \cdots \to \xi_{0,2} \overset{1}{\rightsquigarrow} \xi_{2,0} \to \xi_{3,0} \to \cdots \to \xi_{k,0},
\end{aligned}$$



(I) $q = (\xi^o)$, $k = m$
$2 \leq m < \infty$

(II) $q = (\xi^o)$, $k = 2$
m even

(III) $q = (\xi^o)$, $k = 4$

(IV) $q = (\xi^o, \zeta^o)$, $k = 1$
m odd

(V) $q = (\xi^o, z, \zeta^o)$, $k = 1$

(VI) $q = (\xi^o, z, \zeta^o)$, $k = 2$

(VII) $q = (\xi^o, z, \zeta^o)$, $k = 2$

(VIII) $q = (\xi^o, \zeta^o)$, $k = 2$

(IX) $q = (\xi^o)$, $k = 2$
m even

(X) $q = (\xi^o)$, $k = 4$

(XI) $q = (\xi^o, \zeta^o)$, $k = 2$

Figure 1: The eleven subgraphs Γ_J



$$q = (\xi^o, z) \in \mathcal{F}^I_{\xi^o, z}$$

Figure 2: A sliding move



$$q = (\xi^o, z, \xi^\dagger) \in \mathcal{F}^I_{\xi^o, \xi^\dagger}$$

Figure 3: A switching move

where $\rightarrow$ *and* $\overset{i}{\rightsquigarrow}$ *denote a sliding and a switching at* y_i*, respectively (note that any* $\xi_{i,j} \in \mathcal{E}^I_x$ *admits at most two moves). Any non-backtracking sequence of consecutive moves in* I *from* $\xi_{k,0}$ *to* $\xi_{k,0}$ *is a repetition of the above one (possibly in reverse orientation). This means that, if some nontrivial element of* $\mathcal{F}^I_{y_k}$ *is realized by consecutive moves in* I *from* $\xi_{k,0}$ *to* $\xi_{k,0}$*, then all the moves in the above sequence indeed exist.*

In this case, observe that the above sequence visits all $\xi_{i,j}$ *with* $i-j \not\equiv 0, \pm 1 \pmod{n}$*; so such* $\xi_{i,j}$ *must be an element of* $\mathcal{E}^I_x$*, namely* $m_{i,j} < \infty$*. We also have* $m_{i,j} = 2$ *for these* i, j*, since otherwise* $\xi_{i,j} \in \mathcal{E}^I_x$ *admits no move in* I*, contradicting the above observation. On the other hand, if* $0 \le i, j \le n-1$ *and* $j \equiv i+1 \pmod{n}$*, then we have* $m_{i,j} = 3$*, since the switching* $\xi_{i,\ell} \overset{j}{\rightsquigarrow} \xi_{\ell,i}$ *with* $\ell \equiv j+1 \pmod{n}$*, involved in above sequence, must exist.*

Summarizing, no nontrivial element of $\mathcal{F}^I_{y_k}$ *is realized in this manner unless* Γ_I *is of type* $\widetilde{A}_{n-1}$*. On the other hand, if* Γ_I *is of type* $\widetilde{A}_{n-1}$*, then an element* $Q^\ell \in \mathcal{F}^I_{y_k}$ *(where* Q *is the cycle* $(y_k, y_{k+1}, \dots, y_{n-1}, y_0, \dots, y_k)$*) is realized in this manner if and only if* ℓ *is a multiple of* $n-2$*, since the mover rounds the cycle* $n-2$ *times under the above sequence of moves. Hence in any case, the nontrivial element* $Q \in \mathcal{F}^I_{y_k}$ *is not realized in this manner.*

Example 3.5. *Here we consider the case that* $n = 2N$ *is even (so* $N \ge 2$*) and* $m_{0,k}$ *is even for some* $2 \le k \le n-2$*. Now the shortest non-backtracking nonempty sequence of consecutive moves in* I *from* $\xi_{k,0}$ *to* $\xi_{k,0}$ *(if exists) is*



Figure 4: Coxeter graph in Example 3.5

(up to orientation) the following:

$$\begin{aligned}&\xi_{k,0} \to \xi_{k+1,0} \to \cdots \to \xi_{2N-2,0} \overset{2N-1}{\rightsquigarrow} \xi_{0,2N-2} \to \cdots \to \xi_{2N-4,2N-2}\\ &\quad \overset{2N-3}{\rightsquigarrow} \xi_{2N-2,2N-4} \to \cdots\cdots \to \xi_{0,2} \overset{1}{\rightsquigarrow} \xi_{2,0} \to \xi_{3,0} \to \cdots \to \xi_{k,0},\end{aligned}$$

which visits $\xi_{i,j}$ *with* $i - j \not\equiv 0, \pm 1 \pmod{n}$ if and only if j is even*. Thus it follows in the same way as the previous example that, if some nontrivial element of* $\mathcal{F}^I_{y_k}$ *is realized by consecutive moves in* I *from* $\xi_{k,0}$ *to* $\xi_{k,0}$*, then we have* $m_{i,j} = 2$ *for any* $0 \leq i, j \leq n-1$ *such that* $i - j \not\equiv 0, \pm 1 \pmod{n}$ and either i or j is even*, and* $m_{i,j} = 3$ *for any* i, j *with* $j \equiv i + 1 \pmod{n}$*. Note that the mover rounds the cycle* $N - 1$ *times under the above sequence of moves.*

In this case, if $n \geq 6$*, then it is deduced by applying the above argument to* $\xi_{4,1} \in \mathcal{E}^I_x$ *instead of* $\xi_{k,0}$ *that no nontrivial element of* $\mathcal{F}^I_{y_4}$ *is realized by consecutive moves in* I *from* $\xi_{4,1}$ *to* $\xi_{4,1}$ *unless* Γ_I *is of type* $\widetilde{A}_{n-1}$*. Similarly, if* $n = 4$ *and* $m_{1,3} < \infty$*, then no nontrivial element of* $\mathcal{F}^I_{y_1}$ *is realized by consecutive moves in* I *from* $\xi_{1,3} \in \mathcal{E}^I_x$ *to* $\xi_{1,3}$ *unless* Γ_I *is of type* $\widetilde{A}_3$*.*

Summarizing, by putting $Q = (y_k, y_{k+1}, \dots, y_{n-1}, y_0, \dots, y_k)$*, the nontrivial element* $Q \in \mathcal{F}^I_{y_k}$ *is not realized by consecutive moves in* I *from* $\xi_{k,0}$ *to* $\xi_{k,0}$ *unless* $N = 2$ *(i.e.* $n = 4$*),* $m_{0,2} = 2$ *and* $m_{0,1} = m_{1,2} = m_{2,3} = m_{3,0} = 3$*. On the other hand, if some nontrivial element of* $\mathcal{F}^I_{\xi^o}$ *is realized by consecutive moves in* I *from* ξ *to* ξ *for any* $\xi \in \mathcal{E}^I_x$*, then we have either* Γ_I *is of type* $\widetilde{A}_{n-1}$*, or* $n = 4$*,* $m_{0,2} = 2$*,* $m_{0,1} = m_{1,2} = m_{2,3} = m_{3,0} = 3$ *and* $m_{1,3} = \infty$ *(see Figure 4 for the latter case).*

We close the section by the following two remarks.

Remark 3.6. *Suppose that* $x \in I \subseteq J \subseteq S$*. Then* $\mathcal{E}^I_x \subseteq \mathcal{E}^J_x$ *by definition; while* $\mathcal{F}^I \subseteq \mathcal{F}^J$ *(see Section 2.2) and the isomorphism* π^I *is the restriction of* π^J*. Thus* $\gamma^I_x(\xi; c) = \gamma^J_x(\xi; c)$ *for any* $\xi \in \mathcal{E}^I_x$ *and* $c \in \mathcal{F}^I_{x,\xi^o}$*, so we have* $\Pi^x_I \subseteq \Pi^x_J$ *and* $R^x_I \subseteq R^x_J$*.*

Remark 3.7. *Suppose that* $x \in I \subseteq S$ *and* $y \in I^{\mathrm{odd}}_{\sim x}$*. Then* $\mathcal{E}^I_x = \mathcal{E}^I_y$ *by definition. Moreover, putting* $w = \pi^I(p^I_{y,x}) \in Y^I_{y,x}$*, Theorem 3.2 shows that the* *map* $u \mapsto wuw^{-1}$ *is a bijection* $R_I^x \to R_I^y$ *preserving the order of the product of any two elements; so it gives rise to an isomorphism from* $(W_I^{\perp x}, R_I^x)$ *to* $(W_I^{\perp y}, R_I^y)$ *(cf. [11, Theorem 4.6 (4)]).*

4 Characterization of finitely generated $W_S^{\perp x}$

In this section, we detect completely when the reflection subgroup $W^{\perp x} = W_S^{\perp x}$ introduced in Section 3 is finitely generated (Theorems 4.13 and 4.14). This result suggests that a finitely generated Coxeter group may possess a non-finitely generated reflection subgroup frequently.

4.1 Preliminary steps

First we give some remarks. Let $x \in I \subseteq S$. Since $(W_I^{\perp x}, R_I^x)$ is a Coxeter system, Proposition 2.8 shows that $W_I^{\perp x}$ is finitely generated if and only if $|R_I^x| < \infty$. Moreover, for $y \in I_{\sim x}^{\mathrm{odd}}$ (see Definition 2.10 for notation), Remark 3.7 implies that $|R_I^x| < \infty$ if and only if $|R_I^y| < \infty$. For $x \in J \subseteq I$, Remark 3.6 shows that $|R_J^x| < \infty$ whenever $|R_I^x| < \infty$.

Lemma 4.1. *Let* $x \in I \subseteq S$*, and suppose that* $(\Gamma_I^{\mathrm{odd}})_{\sim x}$ *contains a cycle, and for some* $\xi \in \mathcal{E}_x^I$*,*

$$\text{no } 1 \neq c \in \mathcal{F}_{\xi^o}^I \text{ is realized by consecutive moves in } I \text{ from } \xi \text{ to } \xi. \tag{4.1}$$

Then all the elements $r_x^I(\xi; c) \in R_I^x$ *for* $c \in \mathcal{F}_{x,\xi^o}^I$ *are distinct to each other, and* $W_I^{\perp x}$ *is not finitely generated.*

Proof. The former part of the claim follows immediately from the condition (4.1) and Theorem 3.3. For the other part, the hypothesis on $(\Gamma_I^{\mathrm{odd}})_{\sim x}$ implies that the free group $\mathcal{F}_{\xi^o}^I$ is nontrivial, so $|\mathcal{F}_{x,\xi^o}^I| = |\mathcal{F}_{\xi^o}^I| = \infty$. Thus we have $|R_I^x| = \infty$ by the former claim, as desired. ☐

Here we call a closed path $(x_0, x_1, \dots, x_{n-1}, x_0)$ in a graph *cyclically non-backtracking* if $x_i \neq x_j$ for any $0 \leq i, j \leq n-1$ with $j \equiv i + 2 \pmod{n}$.

Lemma 4.2. *Let* $x \in I \subseteq S$*, and suppose that for some* $\xi \in \mathcal{E}_x^I$*, there is a non-backtracking closed path* $P = (y_0, y_1, \dots, y_{n-1}, y_0)$ *in* Γ_I^{odd} *of positive length such that* $y_0 = \xi^o$*,* $\xi^\dagger \notin V(P)$ *and* $\xi^\dagger$ *is adjacent to* $V(P)$ *in* Γ_I*. Then all the roots* $\gamma_x^I(\xi; p_{x,\xi^o}^I P^k) \in \Pi_I^x$ *for* $k \in \mathbb{Z}$ *are distinct to each other. Hence* $|R_I^x| = |\Pi_I^x| = \infty$*.*

Proof. Note that $\widetilde{\alpha}_{\xi^o,\xi^\dagger} = \lambda\alpha_{\xi^o} + \mu\alpha_{\xi^\dagger}$ for some $\lambda \geq 0$ and $\mu > 0$. Moreover, the non-backtracking closed path P can be decomposed as $P = QCQ^{-1}$, where $Q = (y_0, y_1, \dots, y_\ell)$ for some $\ell \geq 0$ (so $Q^{-1} = (y_\ell, \dots, y_1, y_0)$) and $C = (z_0, z_1, \dots, z_{r-1}, z_0)$ is a cyclically non-backtracking closed path of positive length such that $z_1 \neq y_{\ell-1} \neq z_{r-1}$ if $\ell \geq 1$. Now for $k \in \mathbb{Z} \smallsetminus 0$, we have $P^k = QC^kQ^{-1}$ in $\mathcal{F}^I_{y_0}$, and QC^kQ^{-1} is a non-backtracking closed path of positive length with vertex set $V(P)$. Thus, since $\mathrm{supp}(\pi^I(s,t)) = \{s,t\}$ (see Definition 2.6 for notation) for any edge (s,t) of Γ^{odd}_I, the combination of Lemma 2.7 and Theorem 3.1 implies that

$$\mathrm{supp}(\pi^I(P^k)) = \mathrm{supp}(\pi^I(QC^kQ^{-1})) = V(QC^kQ^{-1}) = V(P),$$

so $\pi^I(P^k) \cdot \alpha_{\xi^\dagger} \neq \alpha_{\xi^\dagger}$ by the hypothesis and Lemma 2.9, while we have $\pi^I(P^k) \cdot \alpha_{\xi^o} = \alpha_{\xi^o}$ since $\pi^I(P^k) \in Y^I_{\xi^o}$. Hence we have $\pi^I(P^k) \cdot \widetilde{\alpha}_{\xi^o,\xi^\dagger} \neq \widetilde{\alpha}_{\xi^o,\xi^\dagger}$; so $\pi^I(P^i) \cdot \widetilde{\alpha}_{\xi^o,\xi^\dagger} \neq \pi^I(P^j) \cdot \widetilde{\alpha}_{\xi^o,\xi^\dagger}$ whenever $i \neq j$. Thus the claim follows, since $\gamma^I_x(\xi; p^I_{x,\xi^o}P^k) = \pi^I(p^I_{x,\xi^o}) \cdot (\pi^I(P^k) \cdot \widetilde{\alpha}_{\xi^o,\xi^\dagger})$. ☐

Now Examples 3.4 and 3.5 yield the following:

Lemma 4.3. *Let $x \in I \subseteq S$ and $J \subseteq I^{\mathrm{odd}}_{\sim x}$, and suppose that Γ^{odd}_J is a cycle in $(\Gamma^{\mathrm{odd}}_I)_{\sim x}$ and $m_{s,t}$ is even for some $s,t \in J$ (so $|J| \geq 4$). If $W^{\perp x}_I$ is finitely generated, then we have either Γ_J is of type $\widetilde{A}_{n-1}$ with $n = |J|$, or $|J| = 4$ and Γ_J is as in Figure 4.*

Proof. Since $J \subseteq I^{\mathrm{odd}}_{\sim x}$, we may assume that $x \in J$ (see Remark 3.7). Then $W^{\perp x}_J$ is also finitely generated, so by Lemma 4.1 (applied to J instead of I), no element $\xi \in \mathcal{E}^J_x$ satisfies the condition (4.1). Since $\mathcal{E}^J_x \neq \emptyset$ by the hypothesis, the claim follows immediately from Examples 3.4 and 3.5. ☐

The next proposition is a key observation in our argument.

Proposition 4.4. *Let $x \in I \subseteq S$ and $J \subseteq I^{\mathrm{odd}}_{\sim x}$ such that Γ^{odd}_J is a cycle in $(\Gamma^{\mathrm{odd}}_I)_{\sim x}$. If $W^{\perp x}_I$ is finitely generated, then for any $\xi \in \mathcal{E}^I_x$ such that $\xi^\dagger \in I^{\mathrm{odd}}_{\sim x}$, we have $m_{\xi^o,\xi^\dagger} = 2$ and $\xi^o, \xi^\dagger \in J$.*

Proof. The claim $m_{\xi^o,\xi^\dagger} = 2$ follows immediately from Lemma 4.1, since ξ satisfies (4.1) whenever $m_{\xi^o,\xi^\dagger} \neq 2$ (now ξ admits no moves in I). So we show that $\xi^o, \xi^\dagger \in J$. Actually, we prove the equivalent claim that $W^{\perp x}_I$ is not finitely generated if $\mathcal{E}^I_x$ contains an element ξ with $m_{\xi^o,\xi^\dagger} = 2$, $\xi^\dagger \in I^{\mathrm{odd}}_{\sim x}$ and $\{\xi^o, \xi^\dagger\} \not\subseteq J$.

Since $(\xi^\dagger, \xi^o)$ is also such an element of $\mathcal{E}^I_x$, we may assume by symmetry that the distance in Γ^{odd}_I from ξ^o to J is less than or equal to that from $\xi^\dagger$ to J (so $\xi^\dagger \notin J$ since $\{\xi^o, \xi^\dagger\} \not\subseteq J$). Let P be one of the shortest paths in Γ_I^{odd} between ξ^o and J, so $\xi^\dagger \not\in V(P)$. If $\xi^\dagger$ is adjacent to $V(P) \cup J$ in Γ_I, then the claim follows from Lemma 4.2 (consider the non-backtracking closed path starting from ξ^o and passing the path P, the cycle Γ_J^{odd} and the path P in this order). So suppose that $\xi^\dagger$ is not adjacent to $V(P) \cup J$. Let $Q = (y_0, y_1, \dots, y_r)$ be one of the shortest paths in Γ_I^{odd} between $y_r = \xi^\dagger$ and some vertex $y_0 \in J$ (so $r \geq 2$ by the above assumption). Then by the above assumption, there is $\zeta \in \mathcal{E}_x^I$ such that $\zeta^o \in J \smallsetminus y_0$, $\zeta^\dagger = y_r$ and $m_{\zeta^o,\zeta^\dagger} = 2$.

Now we prove the claim by induction on the distance r in Γ_I^{odd} between $\zeta^\dagger$ and a cycle Γ_J^{odd} containing ζ^o. The case $r = 1$ has been considered above, so suppose that $r \geq 2$. If $m_{y_1,z}$ is odd for some $z \in J \smallsetminus y_0$, then Γ_I^{odd} contains a cycle which contains ζ^o and y_1 but does not contain $\zeta^\dagger$; so the claim follows from the induction assumption, since the distance between this cycle and $\zeta^\dagger$ is less than r. So suppose not. Now by the choice of Q, the graph Γ_K^{odd} (where $K = J \cup V(Q)$) is the union of the cycle Γ_J^{odd} and the path Q. If $m_{\zeta^o,y_{r-1}} = 2$, then the claim follows from the induction assumption applied to $(\zeta^o, y_{r-1}) \in \mathcal{E}_x^I$. On the other hand, if $m_{\zeta^o,y_{r-1}} \neq 2$, then the element $(y_r, \zeta^o) \in \mathcal{E}_{y_r}^K$ admits no moves in K by the shape of Γ_K^{odd}, so Lemma 4.1 implies that $|R_K^{y_r}| = \infty$ since (y_r, ζ^o) satisfies the condition (4.1). Thus we have $|R_I^x| = |R_I^{y_r}| \geq |R_K^{y_r}| = \infty$, as desired. Hence the proof is concluded. $\square$

From now, let $x \in S$ and put $O = S_{\sim x}^{\mathrm{odd}}$. First, suppose that $W_O^{\perp x}$ is finitely generated, $\mathcal{E}_x^O \neq \emptyset$ and the connected graph Γ_O^{odd} contains a cycle. Take $\xi \in \mathcal{E}_x^O$, and let $\mathcal{P}$ be the family of the subsets $J \subseteq O$ such that Γ_J^{odd} is a single path between ξ^o and $\xi^\dagger$. (Note that $|J| \geq 3$ if $J \in \mathcal{P}$.) Then Proposition 4.4 implies that any cycle in Γ_O^{odd} contains both ξ^o and $\xi^\dagger$, so it is the union of two paths Γ_J^{odd} and $\Gamma_{J'}^{\mathrm{odd}}$ with $J, J' \in \mathcal{P}$. In particular, we have $|\mathcal{P}| \geq 2$ and the union K of all members of $\mathcal{P}$ is a pre-cycle core of Γ_O^{odd} (see Definition 2.1 for terminology). Moreover, if $J, J' \in \mathcal{P}$ and $J \neq J'$, then $\Gamma_{J \cup J'}^{\mathrm{odd}}$ is a single cycle which is the union of Γ_J^{odd} and $\Gamma_{J'}^{\mathrm{odd}}$, since otherwise $\Gamma_{J \cup J'}^{\mathrm{odd}}$ contains a cycle which does not contain ξ^o or $\xi^\dagger$, a contradiction. Thus every vertex in K is contained in some cycle in Γ_O^{odd}, so K is the cycle core of Γ_O^{odd} (see Definition 2.1 for terminology).

Now for distinct $J, J' \in \mathcal{P}$, the group $W_{J \cup J'}^{\perp \xi^o}$ is finitely generated as well as $W_O^{\perp x}$ (see Remarks 3.6 and 3.7), so Lemma 4.3 shows that $\Gamma_{J \cup J'}$ is either of type $\widetilde{A}_{n-1}$ with $n = |J \cup J'| \geq 4$, or as in Figure 4. Moreover, if three $J, J', J'' \in \mathcal{P}$ are distinct, then $\Gamma_{J \cup J'}$ is not of type $\widetilde{A}_{n-1}$ (so is as in Figure 4), since otherwise we have $m_{s,t} = 2$ for some $s \in J \smallsetminus \{\xi^o, \xi^\dagger\}$ and $t \in J' \smallsetminus \{\xi^o, \xi^\dagger\}$ and so the cycle $\Gamma_{J \cup J'}^{\mathrm{odd}}$ and $(s, t) \in \mathcal{E}_x^I$ contradict Proposition 4.4.

Owing to this argument, we have the following:

Proposition 4.5. *Let $x \in S$ and $O = S_{\sim x}^{\mathrm{odd}}$, and suppose that Γ_O^{odd} contains* *a cycle. Let K be the cycle core of Γ_O^{odd} (see Definition 2.1 for terminology). Then $W_O^{\perp x}$ is finitely generated if and only if one of the following three conditions are satisfied:*

1. *No two $s, t \in O$ satisfy that $m_{s,t}$ is even (or equivalently, $\mathcal{E}_x^O = \emptyset$);*
2. *Γ_K is of type $\widetilde{A}_{n-1}$ with $n \geq 4$, and we have $s, t \in K$ whenever $s, t \in O$ and $m_{s,t}$ is even.*
3. *The set K satisfies the following:*

$$\begin{aligned}&4 \leq |K| \leq \infty, \text{ and there are two } y_1, y_2 \in K \text{ such that}\\ &m_{y_1,y_2} = 2,\ m_{y_1,s} = m_{y_2,s} = 3 \text{ for any } s \in K \smallsetminus \{y_1, y_2\}\\ &\text{and } m_{s,t} = \infty \text{ for any distinct } s, t \in K \smallsetminus \{y_1, y_2\},\end{aligned}$$

and $\{s, t\} = \{y_1, y_2\}$ whenever $s, t \in O$ and $m_{s,t}$ is even.

Proof. First, we prove the "only if" part. Suppose that $\mathcal{E}_x^O \neq \emptyset$. Then the argument before this proposition implies that K is as in one of Conditions 2 and 3 (the latter condition is satisfied whenever $|\mathcal{P}| \geq 3$). Moreover, if $s, t \in O$ and $m_{s,t}$ is even, then Proposition 4.4 (applied to $(s, t) \in \mathcal{E}_x^O$) implies that any cycle in Γ_O^{odd} contains both s and t, proving that $s, t \in K$. Hence the claim holds (note that in the case of Condition 3, we have $\{s, t\} = \{y_1, y_2\}$ whenever $s, t \in K$ and $m_{s,t}$ is even).

Secondly, we prove the 'if' part. This is trivial in Case 1 (now $R_O^x = \emptyset$). In Cases 2 and 3, we may assume that $x \in K$. Then we have $\mathcal{E}_x^O = \mathcal{E}_x^K$ by the hypothesis. Moreover, since K is the cycle core of Γ_O^{odd}, we have $\mathcal{F}_{y,z}^O = \mathcal{F}_{y,z}^K$ for all $y, z \in K$ by Lemma 2.5.

Now in Case 2, Examples 3.4 and 3.5 imply that there is a positive integer ℓ such that for any $\xi \in \mathcal{E}_x^K$, the element $C_\xi^\ell \in \mathcal{F}_{\xi^o}^O$, where C_ξ denotes the closed path rounding the cycle Γ_K^{odd} once from ξ^o to ξ^o, is realized by consecutive moves in O from ξ to ξ. This implies that, for any $\xi \in \mathcal{E}_x^K = \mathcal{E}_x^O$, two elements $r_x^O(\xi; p_{x,\xi^o}^O C_\xi^i)$ and $r_x^O(\xi; p_{x,\xi^o}^O C_\xi^j)$ of R_O^x (where $i, j \in \mathbb{Z}$) coincide whenever $i \equiv j \pmod{\ell}$ (see Theorem 3.3). Moreover, we have $\mathcal{F}_{x,\xi^o}^O = \{p_{x,\xi^o}^O C_\xi^k \mid k \in \mathbb{Z}\}$ since $\mathcal{F}_{\xi^o}^O = \mathcal{F}_{\xi^o}^K$ is generated by C_ξ. Thus we have $|R_O^x| \leq \ell \cdot |\mathcal{E}_x^K| < \infty$.

In Case 3, for each $i \in \{1, 2\}$, it follows from the shape of Γ_K^{odd} that $\mathcal{F}_{y_i}^O = \mathcal{F}_{y_i}^K$ is generated by the cycles starting from y_i which is the union of two paths Γ_J^{odd} and $\Gamma_{J'}^{\mathrm{odd}}$ with $J, J' \in \mathcal{P}$. Since all these generators can be realized by certain consecutive moves in K from $(y_i, y_{3-i}) \in \mathcal{E}_x^K$ to (y_i, y_{3-i}), Theorem 3.3 implies that all the elements $r_x^O(y_i, y_{3-i}; c) \in R_O^x$ for $c \in \mathcal{F}_{x,y_i}^O$ coincide with each other. Thus we have $|R_O^x| \leq 2$, since $\mathcal{E}_x^O = \mathcal{E}_x^K = \{(y_1, y_2), (y_2, y_1)\}$.

Hence the claim holds in any case. $\square$

Secondly, we consider the case that Γ_O^{odd} is acyclic. Let $\xi \in \mathcal{E}_x^O$, and take the unique neighbor y_ξ of $\xi^\dagger$ in the graph Γ_O^{odd} lying between ξ^o and $\xi^\dagger$. Then two vertices ξ^o and $\xi^\dagger$ belong to distinct connected components of $\Gamma_{O \smallsetminus y_\xi}^{\mathrm{odd}}$, say $\Gamma_{J_{\xi,o}}^{\mathrm{odd}}$ and $\Gamma_{J_{\xi,\dagger}}^{\mathrm{odd}}$, respectively. Moreover, let $z_{\xi,o}$ and $z_{\xi,\dagger}$ be the unique vertices in $J_{\xi,o}$ and $J_{\xi,\dagger}$, respectively, adjacent to y_ξ in Γ_O^{odd} (so $z_{\xi,\dagger} = \xi^\dagger$). Now we have the following:

Lemma 4.6. *Let $x \in S$ and $O = S_{\sim x}^{\mathrm{odd}}$, and suppose that Γ_O^{odd} is acyclic. Let $\xi \in \mathcal{E}_x^O$. If there is a sequence of consecutive moves in O from ξ to $\zeta \in \mathcal{E}_x^O$, then we have $\zeta^\dagger \in \{z_{\xi,o}, z_{\xi,\dagger}\}$ (where we use the above notations).*

Proof. By the shape of Γ_O^{odd} and definition of the moves, it is deduced inductively that any sliding involved in this sequence before the first switching is of the form $(y, z_{\xi,\dagger}) \to (y', z_{\xi,\dagger})$ with $y, y' \in J_{\xi,o}$; the first switching is the one $(z_{\xi,o}, z_{\xi,\dagger}) \rightsquigarrow (z_{\xi,\dagger}, z_{\xi,o})$ at the vertex y_ξ; any sliding between the first and the second switchings is of the form $(y, z_{\xi,o}) \to (y', z_{\xi,o})$ with $y, y' \in J_{\xi,\dagger}$; and the second switching is the one $(z_{\xi,\dagger}, z_{\xi,o}) \rightsquigarrow (z_{\xi,o}, z_{\xi,\dagger})$ at the vertex y_ξ. The claim is deduced by iterating this argument. $\square$

Owing to this lemma, we have the following:

Proposition 4.7. *Let $x \in S$ and $O = S_{\sim x}^{\mathrm{odd}}$, and suppose that Γ_O^{odd} is acyclic. Then $W_O^{\perp x}$ is finitely generated if and only if there exist only finitely many pairs (s, t) of elements $s, t \in O$ with $m_{s,t}$ even (or equivalently, $|\mathcal{E}_x^O| < \infty$).*

Proof. Note that $\mathcal{F}_{x,\xi^o}^O = \{p_{x,\xi^o}^O\}$ for any $\xi \in \mathcal{E}_x^O$. This implies (by Theorem 3.2 (1)) that $|R_O^x| \leq |\mathcal{E}_x^O|$, so the 'if' part follows immediately.

On the other hand, suppose that the number of such pairs (s, t) is infinite. Then there exist infinitely many elements $\xi_\lambda \in \mathcal{E}_x^O$ for which all the vertices $\xi_\lambda^\dagger$ are distinct. Now by Lemma 4.6, when restricting the equivalence relation $\sim_O$ to the infinite set $\{(\xi_\lambda; p_{x,\xi_\lambda^o}^O)\}_\lambda$, each equivalence class consists of at most two members, so the set decomposes into infinitely many equivalence classes. Thus Theorem 3.2 (1) implies that the subset $\{r_x^O(\xi_\lambda)\}_\lambda$ of R_O^x consists of infinitely many elements, so $|R_O^x| = \infty$. Hence the claim follows. $\square$

We introduce some more notations. Let x and O be as above; here we do not place any assumption for the structure of Γ_O^{odd}. Put

$$\begin{aligned} E &= \{s \in S \smallsetminus O \mid m_{y,s} < \infty \text{ for some } y \in O\}, \\ O_2(s) &= \{y \in O \mid m_{y,s} = 2\} \text{ for any } s \in E. \end{aligned} \tag{4.2}$$

Note that $\mathcal{E}_x^S = \mathcal{E}_x^{O \cup E}$, and $m_{y,s}$ is even or ∞ for any $y \in O$ and $s \in E$.

Remark 4.8. *By definition of the moves, the subset* $\mathcal{E}_x^O$ *of* $\mathcal{E}_x^S$ *is closed under moves in* S*. Moreover, we have* $\mathcal{F}_{x,\xi^o}^S = \mathcal{F}_{x,\xi^o}^O$ *for any* $\xi \in \mathcal{E}_x^S$*, and* $p_{y,z}^S = p_{y,z}^O$ *for any* $y, z \in O$*. This implies that an element* $r_x^S(\xi; c) \in R_S^x$ *lies in* R_O^x *if and only if* $\xi \in \mathcal{E}_x^O$ *(note that* $r_x^S(\xi; c) = r_x^O(\xi; c)$ *if* $\xi \in \mathcal{E}_x^O$*).*

Let $\xi \in \mathcal{E}_x^S \smallsetminus \mathcal{E}_x^O$. Then ξ admits no switching by definition (since $\xi^\dagger \not\in O = S_{\sim\xi^o}^{\mathrm{odd}}$). Moreover, if $m_{\xi^o,\xi^\dagger} \neq 2$, then ξ also admits no sliding by definition. If $m_{\xi^o,\xi^\dagger} = 2$, then any sliding in S from ξ is of the form $\xi \to (y, \xi^\dagger)$ with y a neighbor of ξ^o in the graph $\Gamma_{O_2(\xi^\dagger)}^{\mathrm{odd}}$; so iterating, it follows that the elements of $\mathcal{F}^S$ which can be realized by consecutive moves in S from ξ are the elements of $\mathcal{F}_{\xi^o,y}^{O_2(\xi^\dagger)}$ with $y \in O_2(\xi^\dagger)_{\sim\xi^o}^{\mathrm{odd}}$. This implies the following:

Lemma 4.9. *In the notations, let* $\xi \in \mathcal{E}_x^S \smallsetminus \mathcal{E}_x^O$*,* $\zeta \in \mathcal{E}_x^S$*,* $c \in \mathcal{F}_{x,\xi^o}^S$ *and* $d \in \mathcal{F}_{x,\zeta^o}^S$*.*

1. *Suppose that* $m_{\xi^o,\xi^\dagger} \neq 2$*. Then we have* $(\xi; c) \sim_S (\zeta; d)$ *if and only if* $\xi = \zeta$ *and* $c = d$*.*

2. *Suppose that* $m_{\xi^o,\xi^\dagger} = 2$*. Then we have* $(\xi; c) \sim_S (\zeta; d)$ *if and only if* $\xi^\dagger = \zeta^\dagger$*,* $\zeta^o \in O_2(\xi^\dagger)_{\sim\xi^o}^{\mathrm{odd}}$ *and* $c^{-1}d \in \mathcal{F}_{\xi^o,\zeta^o}^{O_2(\xi^\dagger)}$*.*

Corollary 4.10. *In the notations, we have* $|E| < \infty$ *if* $|R_S^x \smallsetminus R_O^x| < \infty$*.*

Proof. For each $s \in E$, there is some $\xi_s \in \mathcal{E}_x^S \smallsetminus \mathcal{E}_x^O$ with $\xi_s^\dagger = s$ by definition of E. Now Lemma 4.9 implies that all elements $r_x^S(\xi_s) \in R_S^x \smallsetminus R_O^x$ (see Remark 4.8) for $s \in E$ are distinct, proving the claim. □

Now we have the following results:

Proposition 4.11. *In the notations, suppose that* Γ_O^{odd} *contains a cycle. Then we have* $|R_S^x \smallsetminus R_O^x| < \infty$ *if and only if all of the following three conditions are satisfied:*

1. $|E| < \infty$*;*

2. $m_{y,s} \in \{2, \infty\}$ *for any* $y \in O$ *and* $s \in E$*;*

3. *for any* $s \in E$*, the set* $O_2(s)$ *is a pre-cycle core of* Γ_O^{odd} *(see Definition 2.1 for terminology).*

Proof. First we show the "only if" part. Condition 1 follows from Corollary 4.10. Now let $s \in E$ and $y \in O$ such that $m_{y,s}$ is even. If $m_{y,s} \neq 2$, then the element $(y, s) \in \mathcal{E}_x^S \smallsetminus \mathcal{E}_x^O$ satisfies (4.1) with $I = S$, so $|R_S^x \smallsetminus R_O^x| = \infty$ by Lemma 4.1 (since $|\mathcal{F}_{x,y}^S| = \infty$ by the hypothesis on Γ_O^{odd}). This is a contradiction, so Condition 2 follows. Moreover, if $m_{y,s} = 2$ and $(\Gamma^{\mathrm{odd}}_{O_2(s)})_{\sim y}$ does not contain some cycle in Γ^{odd}_O, then by taking a non-backtracking closed path P in Γ^{odd}_O from y to y containing this cycle, Lemma 4.2 implies that all elements $r^S_x(y, s; p^S_{x,y}P^k) \in R^x_S \smallsetminus R^x_O$ for $k \in \mathbb{Z}$ are distinct (note that $V(P) \not\subseteq O_2(s)$ by the choice of the cycle, so s is adjacent to $V(P)$ in Γ_O). This also contradicts the hypothesis $|R^x_S \smallsetminus R^x_O| < \infty$; so for any $y \in O_2(s)$, the connected component $(\Gamma^{\mathrm{odd}}_{O_2(s)})_{\sim y}$ contains all cycles in Γ^{odd}_O. In particular, every connected component of $\Gamma^{\mathrm{odd}}_{O_2(s)}$ contains a common vertex, meaning that $\Gamma^{\mathrm{odd}}_{O_2(s)}$ is connected. Thus Condition 3 follows, proving the "only if" part.

Secondly, we show the 'if' part. By Condition 1, it suffices to show that for each $s \in E$, all the elements $r^S_x(\xi; c) \in R^x_S \smallsetminus R^x_O$ with $\xi^\dagger = s$ coincide with each other (yielding $|R^x_S \smallsetminus R^x_O| \leq |E| < \infty$). So let $\xi, \zeta \in \mathcal{E}^S_x$, $c \in \mathcal{F}^S_{x,\xi^o}$ and $d \in \mathcal{F}^S_{x,\zeta^o}$, and suppose that $\xi^\dagger = \zeta^\dagger = s \in E$. Then $m_{\xi^o,s} = m_{\zeta^o,s} = 2$ by Condition 2. Now Condition 3 and Lemma 2.5 imply that $\Gamma^{\mathrm{odd}}_{O_2(s)}$ is connected and $\mathcal{F}^{O_2(s)}_{\xi^o,\zeta^o} = \mathcal{F}^S_{\xi^o,\zeta^o}$. Thus $(\xi; c) \sim_S (\zeta; d)$ by Lemma 4.9 (2), so the claim follows from Theorem 3.2. Hence the proof is concluded. □

Proposition 4.12. *In the notations, suppose that Γ^{odd}_O is acyclic. Then we have $|R^x_S \smallsetminus R^x_O| < \infty$ if and only if the following three conditions are satisfied:*

1. *$|E| < \infty$;*
2. *for each $s \in E$, there are only finitely many $y \in O$ with $m_{y,s} \notin \{2, \infty\}$;*
3. *for each $s \in E$, the graph $\Gamma^{\mathrm{odd}}_{O_2(s)}$ consists of finitely many connected components.*

Proof. Owing to Corollary 4.10, we may assume without loss of generality that $|E| < \infty$. Since Γ^{odd}_O is acyclic, we have $\mathcal{F}^S_{y,z} = \{p^S_{y,z}\}$ for any $y, z \in O$, and $p^S_{y,z} \in \mathcal{F}^{O_2(s)}_{y,z}$ if $z \in O_2(s)^{\mathrm{odd}}_{\sim y}$. Thus we have $|R^x_S \smallsetminus R^x_O| < \infty$ if and only if for each $s \in E$, there are only finitely many elements $r^S_x(\xi) \in R^x_S$ with $\xi^\dagger = s$ (see Remark 4.8). Now for any distinct $\xi, \zeta \in \mathcal{E}^S_x$ with $\xi^\dagger = \zeta^\dagger = s$, Lemma 4.9 and Theorem 3.2 imply that $r^S_x(\xi) = r^S_x(\zeta)$ if and only if $m_{\xi^o,s} = m_{\zeta^o,s} = 2$ and $\zeta^o \in O_2(s)^{\mathrm{odd}}_{\sim \xi^o}$. This shows that the number of the elements $r^S_x(\xi)$ with $\xi^\dagger = s$ is the sum of the number of $y \in O$ with $m_{y,s} \notin \{2, \infty\}$ and the number of connected components of $\Gamma^{\mathrm{odd}}_{O_2(s)}$. Hence the claim follows. □

4.2 Main theorems

Now we are in a position to give our main theorems. Let $x \in S$ and $O = S^{\mathrm{odd}}_{\sim x}$, and define E and $O_2(s)$ as in (4.2). Moreover, when Γ^{odd}_O contains a cycle, let $K \subseteq O$ denote the cycle core of Γ^{odd}_O (see Definition 2.1 for terminology).

Theorem 4.13. *In the above notations, suppose that Γ_O^{odd} contains a cycle. Then $W^{\perp x}$ is finitely generated if and only if all the following four conditions are satisfied:*

1. *The sets $K \subseteq O$ satisfy one of the three conditions:*
   (a) *$m_{y,z}$ is not even for any $y, z \in O$;*
   (b) *Γ_K is of type $\widetilde{A}_{n-1}$ with $n \geq 4$, and we have $y, z \in K$ whenever $y, z \in O$ and $m_{y,z}$ is even;*
   (c) *The set K satisfies the following:*

   $$\begin{aligned}&4 \leq |K| \leq \infty, \text{ and there are two } y_1, y_2 \in K \text{ such that}\\ &m_{y_1,y_2} = 2,\ m_{y_1,s} = m_{y_2,s} = 3 \text{ for any } s \in K \smallsetminus \{y_1, y_2\}\\ &\text{and } m_{s,t} = \infty \text{ for any distinct } s, t \in K \smallsetminus \{y_1, y_2\},\end{aligned}$$

   and we have $\{y, z\} = \{y_1, y_2\}$ whenever $y, z \in O$ and $m_{y,z}$ is even.
2. *The set E is finite.*
3. *We have $m_{y,s} \in \{2, \infty\}$ for any $y \in O$ and $s \in E$.*
4. *For any $s \in E$, the graph $\Gamma_{O_2(s)}^{\mathrm{odd}}$ is connected and contains K.*

Proof. Note that $|R_S^x| < \infty$ if and only if $|R_O^x| < \infty$ and $|R_S^x \smallsetminus R_O^x| < \infty$. Moreover, since Γ_O^{odd} contains a cycle, a set $I \subseteq O$ is a pre-cycle core of Γ_O^{odd} if and only if Γ_I^{odd} is connected and contains the cycle core of Γ_O^{odd}. Thus the claim is deduced by combining Propositions 4.5 and 4.11. □

Theorem 4.14. *In the above notations, suppose that Γ_O^{odd} is acyclic. Then $W^{\perp x}$ is finitely generated if and only if all the following four conditions are satisfied:*

1. *There exist only finitely many pairs (y, z) of elements $y, z \in O$ such that $m_{y,z}$ is even.*
2. *The set E is finite.*
3. *For each $s \in E$, there exist only finitely many $y \in O$ such that $m_{y,s} \notin \{2, \infty\}$.*
4. *For each $s \in E$, the graph $\Gamma_{O_2(s)}^{\mathrm{odd}}$ consists of finitely many connected components.*

Proof. By the same reason as Theorem 4.13, this is immediately deduced by combining Propositions 4.7 and 4.12. □

By restricting these theorems to some subclasses of Coxeter systems, we obtain the following results, where we continue to use the above notations. First, we consider the finitely generated Coxeter groups.

Corollary 4.15. *Suppose that W is finitely generated. Then $W^{\perp x}$ is finitely generated if and only if either Γ_O^{odd} is acyclic, or Γ_O^{odd} contains a cycle and Conditions 1, 3 and 4 in Theorem 4.13 are satisfied.*

Proof. Since $|S| < \infty$, Condition 2 in Theorem 4.13 and Conditions 1–4 in Theorem 4.14 are automatically satisfied. Thus the claim follows immediately from Theorems 4.13 and 4.14. $\square$

In particular, the reflection subgroup $W^{\perp x}$ is finitely generated whenever W is an affine Coxeter group. This is also deduced by a general theorem of Dyer [5, Theorem 5.1], which shows that any reflection subgroup of an affine Coxeter group is finitely generated.

A Coxeter group W is called *2-spherical* if $m_{s,t} < \infty$ for any $s, t \in S$.

Corollary 4.16. *Suppose that W is 2-spherical and irreducible. Then $W^{\perp x}$ is finitely generated if and only if one of the following three conditions is satisfied:*

1. *$m_{s,t}$ is odd for any $s, t \in S$;*
2. *Γ is of type $\widetilde{A}_{n-1}$ with $4 \leq n = |S| < \infty$;*
3. *Γ_O^{odd} is acyclic and $|S| < \infty$.*

Proof. Note that $S = O \cup E$ by definition of E, since W is 2-spherical. Owing to Theorems 4.13 and 4.14, the 'if' part is easily verified (see Corollary 4.15 for Case 3).

For the "only if" part, first we consider the case that Γ_O^{odd} is acyclic. Our aim here is to show that $|S| < \infty$ (see Condition 3), by using the four properties in Theorem 4.14. Since $S = O \cup E$ as above, owing to Property 2 (in Theorem 4.14), it suffices to show that $|O| < \infty$. Now Property 1 says that $m_{x,y}$ are odd for all but finitely many $y \in O$ (since W is 2-spherical). So if $|O| = \infty$, then x possesses infinitely many neighbors in Γ_O^{odd}; while $m_{y,z}$ is even for any two distinct neighbors y, z of x in Γ_O^{odd}. This contradicts Property 1, so the claim follows in this case.

Secondly, we consider the case that Γ_O^{odd} contains a cycle (covered by Theorem 4.13). Now Property 3 (in Theorem 4.13) implies that $m_{y,s} = 2$ for any $y \in O$ and $s \in E = S \smallsetminus O$; so we have $E = \emptyset$ and $O = S$ since W is irreducible and $O \neq \emptyset$. Moreover, Property 1c cannot hold now; while

Property 1a yields Condition 1 here. Finally, if Property 1b holds, then we have $K = O$ and so Condition 2 is satisfied. Indeed, if $K \neq O$, then $O \smallsetminus K$ contains a vertex y adjacent to some $z \in K$ in Γ_O^{odd}. Now taking $z' \in K \smallsetminus z$, the choice of K implies that $m_{z,z'}$ is not odd, while $m_{z,z'}$ is not even by Property 1b, contradicting that $m_{z,z'} < \infty$. Hence the claim follows. □

A Coxeter group W is called *even* if $m_{s,t}$ is even or ∞ for any distinct $s, t \in S$.

Corollary 4.17. *Suppose that W is even. Then $W^{\perp x}$ is finitely generated if and only if there exist only finitely many $s \in S$ with $m_{x,s} < \infty$.*

Proof. Note that Γ_S^{odd} has no edge (so $O = \{x\}$) since W is even. Now in Theorem 4.14, Conditions 1, 3 and 4 are automatically satisfied; so $W^{\perp x}$ is finitely generated if and only if $|E| < \infty$. Thus the claim follows, since now $E = \{s \in S \smallsetminus x \mid m_{x,s} < \infty\}$. □

A Coxeter group W is called *skew-angled* if $m_{s,t} \neq 2$ for any $s, t \in S$.

Corollary 4.18. *Suppose that W is skew-angled and Γ_O^{odd} contains a cycle. Then $W^{\perp x}$ is finitely generated if and only if $m_{y,z}$ is not even for any $y, z \in O$ and $m_{y,s} = \infty$ for any $y \in O$ and $s \in S \smallsetminus O$ (i.e. W is the free product of W_O and $W_{S \smallsetminus O}$).*

Proof. We use Theorem 4.13. The 'if' part follows immediately from the theorem (note that $E = \emptyset$ now), so we show the "only if" part. Since W is skew-angled, Properties 1b and 1c (in Theorem 4.13) cannot hold, so Property 1a holds. Moreover, Property 3 now implies that $m_{y,s} = \infty$ for any $y \in O$ and $s \in E$; so $E = \emptyset$ by definition of E, therefore $m_{y,s} = \infty$ for any $y \in O$ and $s \in S \smallsetminus O$. Hence the claim follows. □

Finally, we consider the following condition for W; any subset $I \subseteq S$ with at least three elements generates an infinite group. (This is equivalent to the property that a certain complex, called the Davis-Vinberg complex associated to (W, S), has dimension at most two.)

Corollary 4.19. *Suppose that any $I \subseteq S$ with $|I| \geq 3$ generates an infinite group, and Γ_O^{odd} contains a cycle. Then $W^{\perp x}$ is finitely generated if and only if $m_{y,z}$ is not even for any $y, z \in O$ and $m_{y,s} = \infty$ for any $y \in O$ and $s \in S \smallsetminus O$.*

Proof. By the same argument as Corollary 4.18, it suffices to prove the "only if" part by using Theorem 4.13. First, the hypothesis implies that there do not exist three elements $z_1, z_2, z_3 \in O$ such that $m_{z_1,z_2} = m_{z_2,z_3} = 3$ and

$m_{z_1,z_3} = 2$ (i.e. these generate a finite Coxeter group of type A_3; see [9] for the classification of finite Coxeter groups). This shows that Properties 1b and 1c (in Theorem 4.13) cannot hold. Moreover, if $s \in E$, then Property 4 implies that $\Gamma^{\mathrm{odd}}_{O_2(s)}$ contains a cycle (note that $K \neq \emptyset$); in particular, the set $O_2(s)$ contains two distinct vertices y, z with $m_{y,z}$ odd. However, now the set $I = \{s, y, z\}$ contradicts the hypothesis; so we have $E = \emptyset$. Hence the claim follows in the same way as Corollary 4.18. □

References

[1] R. E. Borcherds, *Coxeter groups, Lorentzian lattices, and K3 surfaces*, Int. Math. Res. Not. (1998) 1011–1031.

[2] B. Brink, *On centralizers of reflections in Coxeter groups*, Bull. London Math. Soc. **28** (1996) 465–470.

[3] B. Brink, R. B. Howlett, *Normalizers of parabolic subgroups in Coxeter groups*, Invent. Math. **136** (1999) 323–351.

[4] V. V. Deodhar, *A note on subgroups generated by reflections in Coxeter groups*, Arch. Math. **53** (1989) 543–546.

[5] M. Dyer, *Reflection subgroups of Coxeter systems*, J. Algebra **135** (1990) 57–73.

[6] W. N. Franzsen, R. B. Howlett, *Automorphisms of nearly finite Coxeter groups*, Adv. Geom. **3** (2003) 301–338.

[7] S. R. Gal, *On normal subgroups of Coxeter groups generated by standard parabolic subgroups*, Geom. Dedicata **115** (2005) 65–78.

[8] R. B. Howlett, *Normalizers of parabolic subgroups of reflection groups*, J. London Math. Soc. (2) **21** (1980) 62–80.

[9] J. E. Humphreys, Reflection Groups and Coxeter Groups, Cambridge Univ. Press, 1990.

[10] K. Nuida, *Almost central involutions in split extensions of Coxeter groups by graph automorphisms*, arXiv:math.GR/0512210.

[11] K. Nuida, *On centralizers of parabolic subgroups in Coxeter groups*, arXiv:math.GR/0501061.

Koji NUIDA

Graduate School of Mathematical Sciences, University of Tokyo
3-8-1 Komaba, Meguro-ku, Tokyo, 153-8914, Japan

E-mail: nuida@ms.u-tokyo.ac.jp